



HUMAN AND ENVIRONMENTAL HEALTH EFFECTS OF PLASTIC WASTES DISPOSAL

DR. ARVIND KUMAR ¹ | PRABHASH KUMAR ²

¹ RESEARCH SCHOLAR, DEPARTMENT OF ENVIRONMENTAL SCIENCE, MAGADH UNIVERSITY, BODH GAYA

² RESEARCH SCHOLAR, DEPARTMENT OF ENVIRONMENTAL SCIENCE, MAGADH UNIVERSITY, BODH GAYA

ABSTRACT:

Human population increase and consistent demand for plastics and plastic products are responsible for continuous increase in the production of plastics, generation of plastic waste and its accompanied environmental pollution. In 1907 the invention of Bakelite brought about a revolution in materials by introducing truly synthetic plastic resins into world commerce. By the end of the 20th century, plastics had been found to be persistent pollutants of many environmental niches, from Mount Everest to the bottom of the sea. Whether being mistaken for food by animals, flooding low-lying areas by clogging drainage systems, or simply causing significant aesthetic blight, plastics have attracted increasing attention as a large-scale pollutant.

Dichlorodiphenyldichloroethylene, phenanthrene etc. An estimated 8 million tonnes of plastic is yearly released into the ocean, leading to degradation of marine habitat which eventually affects aquatic organisms. Long term usage and exposure of plastics and plastic products to high temperature can lead to leaching of toxic chemical constituents into food, drinks and water. Indiscriminate disposal of plastics on land and open air burning can lead to the release of toxic chemicals into the air causing public health hazards. This paper also presents recommendations for global prevention and control of plastic wastes.

KEYWORDS:

PLASTIC, OCEAN, POLLUTION, HUMAN, WATER & ENVIRONMENT

INTRODUCTION

Plastics are made up of synthetic organic polymers which are widely used in different applications ranging from water bottles, clothing, food packaging, medical supplies, electronic goods, construction materials, etc. In the last six decades, plastics became an indispensable and versatile product with a wide range of properties, chemical composition and applications. Although, plastic was initially assumed to be harmless and inert, however, many years of plastic disposal into the environment has led to diverse associated problems. Environmental pollution by plastic wastes is now recognized widely to be a major environmental burden, especially in the aquatic environment where there is prolong biophysical breakdown of plastics, detrimental negative effects on wildlife, and limited plastic removal options.

In human occupational and residential environment, plastics made of petrol-based polymer are present in high quantity. At the end-of-life of these plastics, they are usually land-filled together with municipal solid waste. Plastics have several toxic constituents among which are phthalates, poly-fluorinated chemicals, bisphenol A (BPA), brominated flame retardants and antimony trioxide which can reach out to have adverse effects on environmental and public health. Plastics in electronic waste (e-waste) have become a serious global environmental and public health concern due to its large production volume and the presence of inadequate management policies in several countries. plastic hazardous substances from e-wastes can

migrate beyond the processing sites and into the environment.

GLOBAL PRODUCTION OF PLASTICS WASTE

In modern life, plastics are ubiquitous. Its early usage dated back to 1600 B.C., at the time when human hands shaped natural rubber and polymerized into different useful objects. Diverse usage and manufacturing of plastics and plastic products began in 1839 when polystyrene (PS) and vulcanized rubber were discovered. Production of Bakelite which is the first truly synthetic polymer was in 1907 in Belgium, however, by 1930, Bakelite was everywhere, especially in fashion, communication and electrical and automotive industries. It took a decade after this for mass production of plastics to begin and it has constantly expanded ever since.

As at 2008, the annual plastic production was estimated to be 245 million tons globally. At present, single-use packaging is the largest sector, accounting for almost 40% of the overall plastic usage, this is followed by consumer goods, materials for construction, automotive, electrical and agriculture applications at 22%, 20%, 9%, 6% and 3%, respectively.

WORLD PRODUCTION RATE OF PLASTICS

Globally, plastic production was estimated to be 380 million tonnes in 2018. Since 1950 to 2018, plastics of about 6.3 billion tonnes have been produced worldwide, 9% and 12% of which have been recycled and incinerated, respectively. Plastics of about 5 million tonnes are yearly

consumed in UK alone, with only about one-quarter recycled, and the rest land filled. It has been suggested by scientists that by 2050, oceans might contain more plastics than fish in terms of weight. Yearly, approximately 500 billion plastic bags are used out of which an estimated 13 million tonnes ends up in the ocean, killing approximately 100,000 marine lives.

FUTURE PROJECTION OF PRODUCTION OF PLASTIC

Plastic productions have increased in twenty-fold since 1964. Globally, approximately 311 million tonnes of plastics were produced in 2014, expected to double in about 20 year time and possibly quadruple by 2050. International Energy Agency World Energy Outlook in 2015 estimated that, the largest application, plastic packaging (26% of the overall volume), is envisaged to have continuous strong growth, which might double within 15 years, with a possibility of fourfold increase by 2050, to about 318 million tonnes yearly, which is higher than the whole plastic industry today.

TYPES OF PLASTIC

There are different types of plastics based on their constituents and type of materials used in their production.

POLYETHYLENE TEREPHTHALATE (PET)

Polyethylene terephthalate (PET) is a type of plastic which is smooth, transparent and relatively thin. It is also called stomach plastics. PET is commonly used during disposable salad dressing, juice, mouthwash, vegetable oil, cosmetics, soft drinks, margarine and water bottles production, because it is anti-inflammatory and fully liquid. PET is also anti-air, preventing entrance of oxygen into it. Antimony trioxide, an inorganic compound, is used as a catalyst for the production of PET and rubber vulcanization. Plastics made from PET must be prevented from high temperatures so as to prevent the leaching of some toxic additives such as acetaldehyde, antimony and phthalates.

Antimony is a possible human carcinogen. Generally, PET is manufactured for single use only.

HIGH-DENSITY POLYETHYLENE

Worldwide, the most used plastic is polyethylene. High-density polyethylene is a heat-resistant plastic produced from petroleum. It is a major constituent of refrigerators, detergent bottles, toys, milk containers, varieties of plastic grocery bags, etc. No phthalates or BPA is present in high-density polyethylene. High-density polyethylene container is generally considered safe for drink and food because it has no reported health risk even though some studies showed that a long time exposure of the plastics to sunlight can make it harmful.

POLYVINYL CHLORIDE (PVC)

Polyvinyl Chloride (PVC), a type of heat-resistant polymer, is used for packaging fruit juice, cooking oil, etc. PVC is considered highly toxic due to the presence of chemical constituents like heavy metals, dioxins, BPA and

phthalates. Depending on non-plasticization, PVC is flexible due to the presence of phthalates. Phthalates are harmful to humans. The entire PVC life cycle which include the production, usage and disposal are capable of causing severe environmental and public health a risk, hence, its usage has considerably reduced. However, due to cost-effectiveness and versatility, PVC remains very popular in the production of consumer goods. PVC have been reported to cause chronic bronchitis, birth defects, genetic changes, cancer, skin diseases, deafness, vision failure, ulcers, liver dysfunction and indigestion.

LOW-DENSITY POLYETHYLENE

Low-density polyethylene is heat resistant, fragile, flexible and rigid. It is commonly used in packaging of milk, frozen foods and juices. Because the plastic does not have any component that is harmful to human body, its usage is termed safe for beverages and food.

POLYPROPYLENE

Polypropylene, a type of plastics, is strong and semi-transparent. It is heavier and stronger than polyethylene. It is used for packaging medicine, yogurt, ketchup, beverage, etc. Plastics made of polypropylene have no harmful substances and like polyethylene, polypropylene containers are considered safe for humans as packages for food and beverages.

POLYSTYRENE

Polystyrene, a type of petroleum-based plastic, contains benzene which is carcinogenic to humans. Polystyrene is commonly used in the production of insulators and packaging materials. Products from styrene are hazardous to health.

POLYCARBONATE

Polycarbonates are used for packaging consumer goods such as reusable bottles. It contains BPA. Due to exposure to high temperature, BPA can be leached from polycarbonate container into the drink or food stored in them. Because BPA's health risk has been reported in several studies, the usage of polycarbonate plastics have greatly decreased.

SIZE OF PLASTICS: MACRO AND MICROPLASTICS

Size of plastics can be used for their classification, aside the plastic types and their chemical composition. There are two major classifications of plastics at sea:

1) Macro (these are plastics higher 20 mm in diameter) and;

2) Micro (plastics which are less than 5 mm in diameter) plastics. Of these two plastic sizes, the micro plastics are the major pollutants documented for deteriorating the ecosystem. This micro plastic is either produced by design and is called primary micro plastics, or they are formed as a result of degradation of macro plastic called secondary micro plastics.

The major issues in plastic waste cantered around the micro plastics due to an increase difficulty in their

monitoring and a greater effect at the physical and chemical levels on environmental and public health, because of their higher volume-to-surface area ratio. Inadequate waste management and indiscriminate dumping are the major routes of entry of micro plastics into the marine environment. Direct production of micro plastics such as plastic pellets is common, as such are used in fabricating larger items as raw material, however, micro plastics can also be produced through mechanical disintegration of larger plastics or plastic products. This is the case in the breakdown of plastic ropes to finer filaments such as microfibers.

Environmental release of large quantities of micro plastics is in form of cosmetic products and cleaning ingredients such as toothpaste and micro beads in face-wash. Because of the health effects of micro plastics, countries like Canada, USA and others are now phasing out their usage in certain personal care products. Reports of recent research suggest that the detrimental effects of micro plastics especially micro beads, micro plastic fibres and degraded macro plastics in aquatic environment might be higher than that of macro plastics, although studies and legislations to manage plastic pollution are still inadequate.

MANAGEMENT OF PLASTIC WASTES

LANDFILLING

Approximately 10% of household waste is plastics and mostly end up on the landfill. Even though land filling is the commonest waste management conventional approach in many countries, however, scarcity of space for landfills is becoming a major problem. For example, historically, land filling was attractive because it is relatively cheap and simple without necessarily requiring treatment, cleaning or separation. 65% of the overall household waste recoverable plastics were sent to landfill plastic waste land filling is the least favoured waste management option. There is a growing environmental and public health concern about the potential effects of landfills because of the types and quantities of toxic chemicals and their potential for leaching at landfill sites.

Environmental pollution and risks to public health can be reduced if the landfills are well-managed, although there are possibilities of soil and groundwater contamination by disintegrated plastic by products and additives that can persist in the environment on long-term basis.

RECYCLING OF PLASTICS

Reprocessing of recovered plastic scraps or wastes into usable products is called plastic recycling. Most plastics are non-biodegradable in nature, hence, the fundamental work is reduction of waste emissions, effective management and recycling of resulting wastes. Recycling of plastics is a major aspect of the worldwide efforts in minimizing the yearly 8 million tonnes of plastics in the waste stream entering the Earth's ocean. According to Hopewell, et al., plastic recycling terminology is complex due to varieties of recovery activities and recycling. There

are four main categories of recycling which are: primary (which involves the mechanical reprocessing of plastics into a new product with equivalent properties), secondary (which involves the mechanical reprocessing of plastics into a product with lower properties), tertiary (which involves the recovery of the chemical constituents of the plastics) and quaternary (which involves energy recovery from the plastics).

In comparison to the lucrative metal recycling but similar to the low value of glass recycling, recycling of plastics is often more challenging because of low density and low value. Also, there are several technical issues to deal with when recycling plastic. Melting together of different plastic types often cause phase-separation similar to oil and water, and they set in these layers. The resulting phase boundaries is responsible for structural weakness in the final product(s), which has limited the application of this polymer blends. This is the case with polyethylene and polypropylene, which are the two plastics commonly manufactured, and therefore has limited their use for recycling. Of recent, block copolymers has been proposed as a form of macromolecular welding flux or molecular stitches in other to overcome this challenge of phase-separation during plastic recycling.

There can be increase in the percentage of plastics with the possibility of full recycling instead of the large quantity generated as wastes if package good manufacturers reduce their mixing of packaging materials and eliminate contaminants. In view of this, a design guide has been issued by the Association of Plastics Recyclers for recyclability of plastics. There has been an increase in the volume of post-consumer plastics recycled since 1990, although it is still incomparable to other items like corrugated fiber board (approximately 70%) and newspaper (approximately 80%).

LAND POLLUTION

In human occupational and residential environment, plastic products are present in large volume. Pollution by plastics and plastic products can damage and contaminate the terrestrial environment and can be subsequently transferred to the aquatic environment. There is a shortage of data on the volume of plastic wastes on land in comparison to the voluminous data which exist on plastic debris in marine habitat, despite the fact that about 80% of plastic waste present at sea originates from land-related sources. Dumping of plastics on land or land filling plastics leads to a biotic and biotic degradation of the plastics, where plastic additives (e.g. stabilizers, harmful colorant moieties, plasticizers and heavy metals) can leach and eventually percolate into various aspects of the environment, thereby causing soil and water contamination. Reports have shown that micro plastics as well as synthetic polymer fibres are still detectable five years after they have been applied to sewage sludge and soils. Chlorinated plastics are capable of leaching out toxic chemicals into the soil and subsequently seep into the underground water or surrounding aquatic system thereby polluting the ecosystem. Methane, a dangerous

greenhouse gas, which significantly contributes to global warming, is released during microbial biodegradation of plastics.

WATER POLLUTION

Approximately 165 million tonnes of plastic wastes were estimated to be present in the oceans of the world in 2012, while an average of 8 million tonnes of plastics are annually released into the ocean, with about 5 trillion plastic pieces floating on the ocean. Typically, plastics in the oceans can degrade within a year but not completely. During this plastic degradation process, toxic chemicals like polystyrene and BPA can be released into the water causing water pollution. Wastes found in the oceans are made up of approximately 80% plastics. Plastic debris which are floating on the ocean can be rapidly colonized by sea organisms and due to persistence on the ocean surface for a long period of time, this may aid the movement of 'alien' or non-native species. Contaminants from micro plastics are bio available for many marine lives because of their presence in benthic and pelagic ecosystems and their small sizes. Within the marine ecosystem, plastics have been reported to concentrate and sorbs contaminants present in the seawater from different other sources. Examples of such contaminants are persistent organic pollutants like nonylphenol, PCBs, Dichlorodiphenyldichloroethylene (DDE) and phenanthrene, with potential to accumulate in several fold on the plastic debris compared to the surrounding seawater. More than 260 species of marine organisms such as turtles, invertebrates, seabirds, fish and mammals ingested or are entangled in or with plastic debris, leading to reduced movement, feeding, reproductive output, ulcers, lacerations and eventual death.

AIR POLLUTION

Carbon dioxide and methane are released into the air when plastic wastes which were land filled finally decompose. During the decomposition of solid waste in landfills in 2008, an estimated CO₂ equivalent (eqCO₂) volume released into the atmosphere was 20 million tonnes. CO₂ is also released into the atmosphere during the burning of plastics and plastic products, and this CO₂ is capable of trapping radiant heat and hinder it from escaping from the earth causing global warming. Air pollution is one of the major environmental threats to public health, and it is responsible for more than 6 million deaths associated with environmental pollution. Open burning of plastics and plastic products releases pollutants such as heavy metals, dioxins, PCBs and furans which when inhaled can cause health risks especially respiratory disorders. The role of plastics in air pollution in the developing and poor countries of the world cannot be overemphasized, and the impact on the future generations may be massive.

EFFECTS OF PLASTIC WASTES ON ANIMALS

Food supplies for human consumption can be adversely affected if animals are poisoned by toxic constituents from wastes of plastics and plastic products. Indeed, reports of

threat to survival of large marine mammals have been documented due to large amount of plastic wastes entering the world oceans.

Animals are exposed to plastic wastes majorly through ingestion and entanglement, however, ingestion is more frequent than entanglement. Most animals in the oceans mistaking plastic wastes dumped in the ocean for food, thereby ingesting them. Furthermore, entanglement in plastic products like nets can cause harm, damage and even death in marine animals. Reports have shown that more than 260 different species of vertebrate and invertebrate animals ingest plastics or are entangled by plastic or plastic products, with more than 400,000 deaths of marine mammals. Marine pollution by plastic wastes majorly affects sea turtles and other species whose main food are jelly fishes because they often confuse discarded plastic bags for jelly fish. A similar situation is common in sea birds which can confuse micro plastics for cuttlefish or with fishes, which can mistake plastic wastes for their natural prey. Ingestion of plastic wastes is capable of causing obstruction and physical damage to bird's digestive system, reduce the digestive ability of the system leading to starvation, malnutrition and eventually, death.

Many birds, turtles, fishes, seals and other marine animals have died by drowning or suffocation as a result of entanglement in plastic debris. Entanglement has been observed to cause health risks in estimated 243 species of marine lives, often ending in fatalities. Animal entanglement by plastic debris also contributes to death from predators, as the animals are unable to untangle themselves and escape. Coral reefs have been damaged by dragging nets and other plastic products along sea beds. Often times, discarded fishing nets also called "ghost nets" trap marine animals, leading to starvation and death.

HUMAN HEALTH EFFECTS OF PLASTIC ADDITIVES

Different additives are used in the production of plastics and they have been reported to have various detrimental effects on humans.

BISPHENOL A (BPA)

Inner linings of food cans, reusable water bottles, and baby bottles are manufactured using BPA. In 2003, an estimated global output of BPA was greater than 2.2 million metric tonnes annually. As a result of repeated usage of beverage and food containers over a long period of time, BPA molecules can leach from the plastics into the drinks and food. The process of BPA leaching from plastics is accelerated by storing acidic or basic items that can breakdown plastic polymers, exposure of the plastic container to high temperature, and by repeated washing of the plastic container. BPA is an endocrine disruptor which mimics oestrogen in females. Women exposed to BPA have damaged health system such as polycyclic ovarian syndrome, obesity, recurrent miscarriages, endometrial hyperplasia and sterility. BPA alters thyroid hormone axis gene expression, thereby altering its biological functions like metabolism and development. Also, BPA increases thyroid hormone receptor transcriptional corepressor

activity causing a decreasing in thyroid hormone receptor activity. This alteration to thyroid axis causes hypothyroidism. Exposure of children and women of reproductive age to elevated concentration of BPA is of great public health concern because of the higher vulnerability of children and developing foetus to BPA compared to adults exposed to similar concentration. Studies have reported a strong association between the concentration of urinary BPA and liver enzyme abnormalities, cardiovascular disease and type 2 diabetes. Also, BPA associated Neuro-behavioural disorders (e.g. autism), male's abnormal urethra/penile development, female early sexual maturation and increase in hormonally-mediated cancers (e.g. breast and prostate cancers) have been reported

PHTHALATES

Phthalates, also called 1, 2-benzenedicarboxylic acids, consist of a diverse group of diesters of phthalic acid which are produced in large volumes from the 1930s. In industrial applications, particularly in the manufacture of food packaging, raincoats, medical devices, toys, hoses, vinyl flooring and shower curtains, high molecular weight phthalates (e.g. di(2-ethylhexyl) phthalate (DEHP)) are commonly used. Phthalates with low molecular weight especially dibutyl phthalate (DBP) and diethyl phthalate (DEP) are used as solvents in the manufacture of products such as lacquers, coatings, varnishes and personal-care products (e.g. cosmetics, perfumes and lotions). Lack of chemical bond between phthalates and the plastic matrix makes it easy for phthalates to reach out and contaminate the environment. Due to the presence of phthalates in many consumer goods, there is widespread human exposure to phthalate.

Phthalates are endocrine disruptors with anti-androgenic activity. Children and infants are mostly exposed to phthalates because of their frequent mouthing of objects like plastic toys and fingers, and direct skin contact with phthalate contaminated substances. Ingestion of phthalates in breast milk, cow milk, or food packaging materials are the main routes of exposure in breast feeding infants. Using personal care products frequently can increase the rate of exposure to phthalates of low molecular weight, indeed, report have shown that men who recently used aftershave and cologne have increased phthalate exposure, while infants that used certain infant-care products such as shampoos, lotions and powders also showed increased exposure. High phthalate concentration alters hormone levels thus causing birth defects in rodents exposed to certain types of phthalates. Butyl benzyl phthalate have been reported to cause rhinitis and eczema in children and has been classified as possible Class-Chuman carcinogen in the 1986 US EPA guidelines.

POLYCHLORINATED BIPHENYLS (PCBS)

Marine food web has been continuously polluted with polychlorinated biphenyls (PCBs) for the last 70years, particularly in seabirds. PCBs ingestion may cause

reproductive disorders, enhance disease proliferation, alters hormone levels and death. PCBs can contaminate marine food web through the plastic bits and it has been shown that PCB is detrimental to marine life even at very low concentrations.

PLASTIC WASTE MANAGEMENT AND RECYCLING

In reducing toxic effects of plastic wastes on the environment and public health, waste management plays a major role. For global reduction of plastic litters and ocean pollution, there is need for improvement in proper plastic waste collection, treatment and disposal. Inadequate management of landfills will make way for harmful chemicals in plastic wastes to leach into the environment, polluting the soil, air and underground water.

Proper wastewater management will prevent micro plastics from entering the environment from the landfills. Most treated wastewaters are discharged into rivers or oceans; therefore, there is need for a ban such as Annex V to the International Convention for Prevention of Pollution from Ship (MARPOL) agreement, which will prevent plastic waste disposal into the sea.

EDUCATION AND PUBLIC AWARENESS

Efforts must be made to educate the general populace on the potential environmental and public health effect of pollution by plastic wastes. This will go a long way to reduce the pollution rate and preserve the quality of the environment. There is need for people to be aware of the chemical constituents of plastic products and their health effects. Educational curriculums at different levels must include ways of plastic pollution reduction and waste management systems as information resources.

BIOPLASTICS AS ALTERNATIVE

Bioplastics is a plastic produced from cellulose that is made of wood pulp by a British chemist in the 1850s. Now, bioplastics can be produced from different biodegradable and non-biodegradable materials including weeds, hemp, plant oil, potato starch, cellulose, corn starch, etc.. Sugar-based bioplastics can biodegrade under normal conditions for composting. Bioplastics are environmentally friendly since they require less fossil fuels during production in comparison to other types of plastic.

Although bioplastics have been used commercially in just few applications, they are widely used in consumer goods for items that are disposable like cutlery, bowls, pots, crockery, straws and packaging. In principle, bioplastics can replace petroleum-derived plastics in many applications, however, the problem lies with the cost and performance of bioplastics. If there are no specific regulations globally to limit the use of conventional plastics, there may be no favourable usage of bioplastics. For example, Italy has since 2011 enacted law that made it compulsory for biodegradable plastic bags to be used for shopping. In the production of bioplastics, substitute for fossil fuel resources like wood, cellulose, sugar and starch are used. This has made bioplastic production more sustainable and environmentally friendly in comparison to

conventional plastic production. The production of bioplastics decreases consumption of non-renewable energy and reduces the emission of greenhouse gases.

We believed that the problem of plastic waste generation and the accompanied environmental and public health effects can be handled if globally, manufacturers can embrace the use of bioplastics. The biodegradability with little or no toxic products left behind will go a long way to protect our natural environment from the menace of conventional plastic wastes, protect our world's organisms and make the world safe for humans.

REFERENCES

1. Nagy A, Kuti R (2016) The Environmental Impact of plastic waste Incineration. AARMS 15: 231-237.
2. Foldi L, Halasz L (2009) Környezetbiztonság. Complex KiadóKft, Budapest, Hungary.
3. Padanyi J, Foldi L (2014) Environmental responsibilities of the military soldiers have to be "Greener Berets". Ecology Manage 2: 48-55.
4. Hardesty BD, Good TP, Wilcox C (2015) Novel methods, new results and science based solutions to tackle marine debris impacts on wildlife. Ocean & Coastal Management 115: 4-9.
5. Creton C (2017) Molecular stitches for enhanced recycling of packaging. Science 355: 797-798.
6. Eagan JM, Xu J, Di Girolamo R, Thurber CM, Macosko CW, et al. (2017) Combining polyethylene and polypropylene: Enhanced performance with PE/iPP multiblock polymers. Science 355: 814-816.
7. Fleischman T (2018) Polymer additive could revolutionize plastics recycling. Cornell. Edu. Cornell University.
8. Alan Bridgewater, Gill Bridgewater (2007) The Self-Sufficiency Handbook: A Complete Guide to Greener Living. Skyhorse Publishing Inc., USA, 62.
9. Public and Environmental Health Effects of Plastic Wastes Disposal: (2019) Okunola A Alabi.
10. Mathalon, A., Hill, P., 2014. Microplastic fibers in the intertidal ecosystem surrounding Halifax Harbor, Nova Scotia. Mar Pollut Bull 81 (1), 69-79.
11. Shobha ray, sherghati nishu, pollution create by family, start 2013- end 17 April 2022 Book. Pg- 13(10)1989, hotspot.
12. Li, J., Qu, X., Su, L., Zhang, W., Yang, D., Kolandhasamy, P., Li, D., Shi, H., 2016a. Microplastics in mussels along the coastal waters of China. Environ Pollut 214, 177-184.
13. Qu, X., Su, L., Li, H., Liang, M., Shi, H., 2018. Assessing the relationship between the abundance and properties of microplastics in water and in mussels. Sci Total Environ 621, 679-686.
14. Zeytin, S., Wagner, G., Mackay-Roberts, N., Gerdts, G., Schuirmann, E., Klockmann, S., Slater, M., 2020. Quantifying microplastic translocation from feed to the fillet in European sea bass *Dicentrarchus labrax*. Mar Pollut Bull 156, 111210.
15. Karami, A., Golieskardi, A., Choo, C. K., Larat, V., Karbalaee, S., Salamatinia, B., 2018. Microplastic and mesoplastic contamination in canned sardines and sprats. Sci Total Environ 612, 1380-1386.
16. Lamb, J. B., Willis, B. L., Fiorenza, E. A., Couch, C. S., Howard, R., Rader, D. N., True, J. D., Kelly, L. A., Ahmad, A., Jompa, J., Harvell, C. D., 2018. Plastic waste associated with disease on coral reefs. Science 359 (6374), 460-462.
17. Al-Jufaili, S., Al-Jabri, M., Al-Baluchi, A., Baldwin, R. M., Wilson, S. C., West, F., Matthews, A. D., 1999. Human Impacts on Coral Reefs in the Sultanate of Oman. Estuar Coast Shelf Sci 49, 65-74.
18. Angiolillo, M., Lorenzo, B. D., Farcomeni, A., Bo, M., Bavestrello, G., Santangelo, G., Cau, A., Mastas-cusa, V., Cau, A., Sacco, F., Canese, S., 2015. Distribution and assessment of marine debris in the deep Tyrrhenian Sea (NW Mediterranean Sea, Italy). Mar Pollut Bull 92 (1-2), 149-159.
19. Rochman CM, Hoh E, Kurobe T, The SJ (2013b) Ingested plastic transfers hazardous chemicals to fish and induces hepatic stress. Sci Rep 3: 3263.
20. Comanita ED, Hlihor RM, Ghinea C, Gavrilescu M (2016) Occurrence of plastic waste in the environment: Ecological and Health risks. Environ Engineer Manag J 15: 675-685.
21. Mouat J, Lozano RL, Bateson H (2010) Economic Impacts of marine litter. KIMO Int conference 105.